



## CHEMICAL COMPOSITION OF PRECIPITATION AT TWO SAMPLING SITES IN MEXICO: A 7-YEAR STUDY

A. P. BAEZ, R. D. BELMONT and H. G. PADILLA

Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México, Ciudad Universitaria,  
México, D.F., 04510, Mexico

(First received 5 April 1995 and in final form 1 May 1996)

**Abstract**—This paper describes the results of a study of the chemical composition of bulk and wet precipitation, collected for 7 years in a large urban area and a forested rural site in Central Mexico. In rain samples from both sites, sulfate was the most abundant ion followed by ammonium. Seasonal patterns of monthly volume-weighted concentration show that the highest values of  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{Ca}^{2+}$  and  $\text{NH}_4^+$  were measured in May, the beginning of the rainy season in Central Mexico. The lowest values were observed from July to September, the months when most thunderstorms occurred. Multiple regression correlation analysis was applied to all data and a hierarchical model was used to select the variables included in the final model. In Rancho Viejo the model showed that  $\text{SO}_4^{2-}$  contributed with 41% in the  $\text{H}^+$  concentration prediction. In Mexico City, despite the high  $\text{SO}_4^{2-}$  concentration in rain water, this ion only contributed 0.46% whereas  $\text{Ca}^{2+}$  contributed in wet and bulk precipitation, 61 and 41%, respectively. Finally no long-term trends were noticed. Copyright © 1996 Elsevier Science Ltd

*Key word index:* Rain chemistry, urban-rural comparison.

### INTRODUCTION

Many studies have been made on the chemical composition of precipitation over extended intervals, to detect modifications caused by the increase of human activities, but it has been difficult to show trends due to the wide variations observed in rain water composition (Colin *et al.*, 1990). Precipitation, however, has unique scavenging properties, which makes it a useful indicator of ambient pollution levels. Its analysis yields valuable data concerning inputs of both nutrients and toxic compounds to terrestrial and aquatic systems (Caiazza *et al.*, 1978; Peden and Skowron, 1978; Kins, 1982; Pratt *et al.*, 1983; Baez and Belmont, 1987; Parungo *et al.*, 1990).

Wet deposition for a given area depends on total precipitation and on factors that determine concentration variations of the chemical species of interest in rainfall (Prado-Fiedler, 1990).

In this study we analyzed the composition of daily precipitation during a 7-year period, its annual variation, and determined an annual estimate of wet and bulk precipitation. We also determined the precipitation chemistry in a highly polluted urban area and in a forested rural region.

### EXPERIMENTAL METHODS

#### *Sampling sites*

Two sampling sites were selected: an urban area and a forested rural region. The selected urban area was Mexico

City. The collector was located on the roof of the Atmospheric Sciences Center's building, at the campus of the National University, at an elevation of 2250 m above sea level (ASL). The mean annual temperature is 16°C and the mean annual rainfall is 860 mm at this location. Prevailing winds during the rainy season come from the southeast to northeast and usually the clouds are of the convective type (Jauregui, 1986). The second sampling site (Rancho Viejo) was located in a forested region, 80 km southwest of Mexico City, in the State of Mexico, at 2700 m (ASL) with a mean annual rainfall of 1440 mm. During the rainy season, easterly winds at 700–600 mb (Servicio Meteorológico Nacional, 1992) transport air pollutants, emitted in Mexico City, to Rancho Viejo, according to the synoptic conditions that prevail during summer. The locations of the sampling sites are shown in Fig. 1.

#### *Sampling methods*

From 1987 to 1990, bulk precipitation was collected daily at the Mexico City site using a polyethylene funnel 24 cm in diameter, draining into a 3 l polyethylene bottle inside a sealed polyurethane insulated box to reduce the evaporation and growth of microorganisms. The top of the funnel was placed 1.80 m above ground level to prevent contamination from splashing. Funnel and bottles were rinsed thoroughly with deionized water before exposure. From 1991 to 1993, only wet precipitation samples were collected in Mexico City, on a daily basis, with an automatic wet/dry precipitation collector (Andersen, General Metal Works, Inc.) into a standard high-density polyethylene bucket. After collection, bottles were sealed with polyethylene caps and the buckets were tightly sealed with polyethylene covers. In both cases, bottles and buckets were sent to the Atmospheric Sciences Center's Chemical Laboratories in Mexico City for chemical analysis.

At Rancho Viejo only wet precipitation samples were collected from 1987 to 1993. The collection system consisted




Fig. 1. Monitoring network sampling stations: (1) Xalostoc; (2) Pedregal and University of Mexico (MCMA: Mexico City metropolitan area).

of a 24 cm diameter polyethylene funnel, which was 1.80 m above grass-covered ground. The funnel was attached to a 30 cm long polyethylene tubing inserted into a 3.5  $\ell$  polyethylene bottle. The funnel was covered between rain events. The funnel and tubing were rinsed with deionized water shortly before the rain started.

#### Chemical analysis

The samples were analyzed for pH,  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , and  $\text{NH}_4^+$ . For chemical analysis, unfiltered samples were used from 1987 to 1990. From 1991 to 1993, samples were filtered through a 0.45  $\mu\text{m}$  membrane Millipore leached with deionized water before chemical analysis. pH was measured within 24 h upon the arrival of samples to the laboratory. Sulfate,  $\text{NH}_4^+$  and  $\text{NO}_3^-$  were determined by turbidimetric, phenate and salicylate methods, respectively (APHA, 1985; Mészáros, 1978). Since 1992,  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  were determined by ion chromatography (Perkin-Elmer-Isocratic LC pump 250 with a conductivity detector). Calcium and  $\text{Mg}^{2+}$  were measured by flame atomic absorption spectrophotometry (Perkin-Elmer 460). The detection limits (DLs) for  $\text{SO}_4^{2-}$ , turbidimetric and ion chromatography methods, were 1 and 0.2  $\text{mg } \ell^{-1}$ , respectively. The DLs for  $\text{NO}_3^-$ , salicylate and ion chromatography methods, were 0.1 and 0.09  $\text{mg } \ell^{-1}$ , respectively. The detection limit (DL) for  $\text{NH}_4^+$  was 0.013  $\text{mg } \ell^{-1}$ . The DL for  $\text{Ca}^{2+}$  was 0.01  $\text{mg } \ell^{-1}$  and for  $\text{Mg}^{2+}$  it was 0.002  $\text{mg } \ell^{-1}$ .

#### Precision of analysis

Precision was determined by analysis of standards from 10 replicate measurements at different concentrations. The

results showed that the precision for  $\text{SO}_4^{2-}$ ,  $\text{NH}_4^+$  and  $\text{NO}_3^-$  (turbidimetric, phenate and salicylate methods) was < 10% relative standard deviation (RSD) and for  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$  (ion chromatography method),  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  it was < 5% RSD.

#### RESULTS AND DISCUSSION

This study corresponded only to the rainy season in Central Mexico (May to October). For the statistical analysis, whenever a value of a particular ion was below the detection limit, half the detection limit was used for computation purposes.

In the statistical analysis the first step was to plot frequency distributions. Figure 2 shows the frequency distribution for the  $\text{SO}_4^{2-}$  in Mexico City (wet precipitation, 1991–1993) and Rancho Viejo (only wet precipitation, 1987–1993), which follows approximately a log-normal distribution, but skewed toward higher concentration values. The distribution shown in Fig. 2 is similar for the concentrations of most other ionic species in precipitation samples from Mexico City and Rancho Viejo.

Geometric means, volume-weighted means (VWM), standard deviations of the VWM, and minimum and maximum values for the entire data period 1987–1993




Fig. 2. Frequency distribution plots of  $\text{SO}_4^{2-}$  for both sampling sites, which follow an approximately log-normal distribution, but skewed toward higher concentration values.

for all analyzed ions are shown in Table 1. The standard deviations of the VWM were calculated using the formula of Galloway *et al.* (1984).

Figure 3 shows monthly volume-weighted concentrations of rain samples in Mexico City (bulk and wet precipitation) and in Rancho Viejo (only wet precipitation) averaged over the 7-year period. The  $\text{SO}_4^{2-}$

monthly concentration in Mexico City was highest in May, while at Rancho Viejo the highest concentration was observed in June. The lowest concentration for this ion was found from July to September at both the sampling sites. The monthly behavior of  $\text{NO}_3^-$  and alkaline ions  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{NH}_4^+$  followed similar patterns as those of  $\text{SO}_4^{2-}$ . The high values found in

Table 1. Volume-weighted means (VWM), standard deviations of the volume-weighted means (SVWM), geometric means, minima and maxima, in  $\mu\text{eq l}^{-1}$ , for entire data 1987–1993

	$\text{SO}_4^{2-}$	$\text{NO}_3^-$	$\text{Ca}^{2+}$	$\text{Mg}^{2+}$	$\text{NH}_4^+$	$\text{H}^+$
<i>University of Mexico, 1987–1990 (N = 104)</i>						
VWM	131.3	41.45	70.51	9.67	93.51	23.51
SVWM	9.28	3.79	4.57	0.70	7.20	4.28
G. mean	132.6	36.71	70.87	9.28	88.35	5.41
Minimum	34.17	5.29	3.50	0.82	15.71	0.04
Maximum	600	255	344	47.54	498	251
<i>University of Mexico, 1991–1993 (N = 79)</i>						
VWM	75.30	34.82	33.07	3.67	76.16	22.29
SVWM	6.41	3.29	2.82	0.30	5.01	4.00
G. mean	74.61	32.81	31.48	3.63	74.25	4.43
Minimum	17.70	5.50	4.45	0.57	15.00	0.10
Maximum	268	202	363	31.15	264	275
<i>Rancho Viejo, 1987–1993 (N = 267)</i>						
VWM	39.75	15.65	9.26	2.53	26.91	20.50
SVWM	3.17	1.27	0.91	0.32	2.06	1.81
G. mean	32.52	14.19	6.14	1.63	21.96	14.94
Minimum	1.46	1.21	0.50	0.16	0.71	0.11
Maximum	284	179	146	45.90	307	199

N = number of samples.

May for these ions are associated with the beginning of the rainy season. The first rains scavenge the accumulated pollutants resulting in relatively high ionic concentrations.

The increase in  $\text{Ca}^{2+}$  concentration, observed also in May, reflects the increased amounts of soil particles being blown into the atmosphere by wind action (Galloway and Likens, 1978) and dust from lime and cement plants. The washout of large particulate matter takes place during the early stages of rainfall; this particulate matter settles rapidly due to dry fall out, increasing the  $\text{Ca}^{2+}$  content in bulk precipitation more than in wet precipitation (Fig. 3). Saylor *et al.* (1992) have suggested that larger ionic concentrations are due to in-cloud and below-cloud removal of aerosols during brief showers that occur after extended periods of dryness and air stagnation.

In Mexico City during winter and early spring, the atmosphere is characterized by almost daily temperature inversions and stagnation conditions. These conditions contribute to the accumulation of air pollutants emitted in huge amounts by small, medium and large industries located in and near Mexico City, and by more than three million motor vehicles. Therefore, it is reasonable to assume that the highest ionic concentrations occur during the first rains in May or after periods of dryness, as Saylor *et al.* (1992) have also reported. July through September are the months when most thunderstorm activity occurs, easterly winds are well established, and the amounts of precipitation are the highest, resulting in dilution of ionic species.

The annual variations of volume-weighted means for  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  and  $\text{H}^+$  are shown in Fig. 4.

Annual trends were not evaluated, because the data sets showed small annual term variations. This may be due to the high variability of annual inputs of the different ionic species, or to the variation of various meteorological factors such as wind direction (trajectories of the precipitation air masses), rainfall rate, total amounts of rainfall, and height of cloud base (Asman *et al.*, 1981). Care must be taken, however, when interpreting these results since analytical methods were not the same throughout the 7-year sampling period.

To test for significant differences among ionic concentrations in wet precipitation between both sampling sites, the Student's *t*-test (two-tail at 5%) was applied. The results of this statistical test indicate that there were significant differences between these two sampling sites. These variations in ionic concentrations possibly include differences in the meteorological conditions and rain chemistry between events and between locations (Pratt *et al.*, 1983). Additionally, changes in air pollutant concentrations in the atmosphere of Mexico City are important in explaining the variability of rain water chemical composition. Table 2 shows the ion pair  $\log_{10}$  correlation from both sampling sites. For the University of Mexico, the highest correlation coefficients corresponded to the pair  $\text{Ca}^{2+}$ – $\text{Mg}^{2+}$  in both sampling periods. This can be attributed to suspended soil dusts (calcareous soils are predominant in this area) and industrial alkaline particle emissions. The high correlation between  $\text{SO}_4^{2-}$ – $\text{NO}_3^-$ , in both sampling periods, is probably due to the coemission of their precursors  $\text{SO}_2$  and  $\text{NO}_x$ , which are emitted in great amounts by the multiple fossil fuel combustion sources in the Mexico City




Fig. 3. Monthly volume-weighted mean concentrations, in  $\mu\text{eq l}^{-1}$ , of selected ionic components in Mexico City for bulk precipitation and wet precipitation collected at the University of Mexico during 1987–1990 and 1991–1993, respectively, and wet only precipitation collected at Rancho Viejo during the period 1987–1993.

metropolitan area. At Rancho Viejo the observed high correlations are also probably due to the precursors  $\text{SO}_2$  and  $\text{NO}_x$ , which are transported from Mexico City and the industrial area of Toluca. This assumption is supported by the radiosonde observations made at the Mexico City International Airport and wind data from pilot balloons launched in Rancho Viejo, which showed similar wind directions ranging from north–northeast to south–southwest (Padilla *et al.*, 1993). To support the assumption that

$\text{SO}_2$  and  $\text{NO}_x$  are important precursors of  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  due to the coemission of these precursors, a simple linear hourly correlation was calculated between  $\text{NO}_3^-/\text{NO}_x$  and  $\text{SO}_4^{2-}/\text{SO}_2$ , for the period of 1991–1993. The values of  $\text{NO}_x$  and  $\text{SO}_2$  were provided by the Mexico City air quality monitoring network, from two selected sampling stations; one in the northern part of the city (Xalostoc) and one in the southern zone (Pedregal) (Fig. 1). The best correlations found at Xalostoc, between  $\text{NO}_3^-/\text{NO}_x$  were at




Fig. 4. Annual variations of volume-weighted mean concentrations in  $\mu\text{eq l}^{-1}$  for the ionic species  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  and  $\text{H}^+$ , whose trends could not be evaluated because the data showed small variations.

1:00, 11:00, 12:00, 14:00, 19:00 h (local time) and from 21:00 to 24:00 h, and the correlation coefficient ( $r$ ) ranged from 0.399 to 0.624 at  $p < 0.01$ . Between 14:00 and 20:00 h,  $r$  was 0.409 and 0.388, respectively, at  $p < 0.05$ . At Pedregal only the correlation values were found at 1:00, 20:00 and from 22:00 to 24:00 h with  $r$  ranging from 0.314 to 0.514 at  $p < 0.01$ . With respect to  $\text{SO}_2$ , the best correlation between  $\text{SO}_4^{2-}/\text{SO}_2$  was found at the Pedregal sampling station

for all hours with the exception of 1:00 and 24:00 h;  $r$  ranged from 0.361 to 0.433 at  $p < 0.01$ , and  $r$  ranged from 0.466 to 0.488 at  $p < 0.05$ . With respect to the Xalostoc station the only correlation between  $\text{SO}_4^{2-}/\text{SO}_2$  was found at 7:00 h, with  $r = 0.470$  at  $p < 0.01$ , and at 10:00, 17:00, 18:00, 20:00 and 21:00 h,  $r$  ranging from 0.363 to 0.470 at  $p < 0.05$ .

Hydrogen peroxide is an important oxidant of dissolved S(IV) in cloud water and precipitation

(Penkett *et al.*, 1979; Schwartz, 1984; Calvert *et al.*, 1985; Seinfeld, 1986). In addition, observational evidence for the in-cloud reaction of SO<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> has been reported from field studies (Daum *et al.*, 1984). However, the solubility of SO<sub>2</sub> in the liquid phase is strongly pH-dependent, since it is governed by the acid-base equilibria; under acidic conditions the solubility of SO<sub>2</sub> is very low, while it is strongly increased at higher pH (Seinfeld, 1986). Thus, the aqueous concentration of S(IV) depends on the presence of acidic and basic components that establish pH in atmospheric water (Seinfeld, 1986). On the other hand, at pH > 5 the oxidation by ozone is more efficient than by H<sub>2</sub>O<sub>2</sub> (Penkett *et al.*, 1979; Seinfeld, 1986). Measurements of H<sub>2</sub>O<sub>2</sub> were not made during this study; for this reason it is not possible to ascertain the

contribution of H<sub>2</sub>O<sub>2</sub> to the oxidation of the SO<sub>2</sub> and NO<sub>x</sub>, despite the fact that the average pH of the rain water was less than 5. However, we must consider the inputs of NH<sub>4</sub><sup>+</sup>, Ca<sup>2+</sup> and Mg<sup>2+</sup> to the neutralization of the acidity of rain water, which in many cases had a pH > 7. In Mexico City the air quality standard for ozone (0.11 ppm) is exceeded on more than 80% of the days per year; thus, it is possible to assume that there are substantial amounts of ozone for the oxidation of the SO<sub>2</sub> and NO<sub>x</sub> that account for the presence of SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> in the precipitation.

A multiple regression correlation (MRC) analysis was performed on the rain chemistry data. The data followed an approximately log-normal distribution; therefore, a log<sub>10</sub> transformation was applied to all data and the hierarchical model (Cohen and Cohen, 1975) was used to select the variables included in the final model. Table 3 shows the model with the best-fit coefficients for Mexico City (1987–1990 and 1991–1993 periods) and for Rancho Viejo (1987–1993 period).

At Rancho Viejo, the model shows that SO<sub>4</sub><sup>2-</sup> contributed by far the most (41%) to the H<sup>+</sup> concentration prediction. This result is expected from the geographic characteristics of the location, which is downwind from Mexico City with neither nearby local industry nor significant soil particle emission sources.

On the other hand, Ca<sup>2+</sup> contributes with 9%, while Mg<sup>2+</sup>, NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>, combined, account for approximately 7%.

Despite the high SO<sub>4</sub><sup>2-</sup> concentrations found in bulk precipitation collected in Mexico City from 1987 to 1990, this ion is not important in the regression equation. In fact, calcium is the most relevant and important parameter, with a contribution of 38%, whereas sulfate contributes as little as 0.46%. Note the contrast with Rancho Viejo where Ca<sup>2+</sup> and SO<sub>4</sub><sup>2-</sup> contribute 9 and 41%, respectively. We interpret this to indicate that rain chemistry in Mexico

Table 2. Simple ion-pair log<sub>10</sub> correlation coefficients (*r*)

	NO <sub>3</sub> <sup>-</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	NH <sub>4</sub> <sup>+</sup>	H <sup>+</sup>
<i>University of Mexico, 1987–1990 (N = 104)</i>					
SO <sub>4</sub> <sup>2-</sup>	0.760 <sup>a</sup>	0.738 <sup>a</sup>	0.664	0.789 <sup>a</sup>	-0.068 <sup>a</sup>
NO <sub>3</sub> <sup>-</sup>		0.575 <sup>a</sup>	0.551 <sup>a</sup>	0.621 <sup>a</sup>	0.163
Ca <sup>2+</sup>			0.810 <sup>a</sup>	0.507 <sup>a</sup>	-0.458 <sup>a</sup>
Mg <sup>2+</sup>				0.515 <sup>a</sup>	-0.398
NH <sub>4</sub> <sup>+</sup>					0.036
<i>University of Mexico, 1991–1993 (N = 79)</i>					
SO <sub>4</sub> <sup>2-</sup>	0.840 <sup>a</sup>	0.655 <sup>a</sup>	0.702 <sup>a</sup>	0.778 <sup>a</sup>	0.234 <sup>b</sup>
NO <sub>3</sub> <sup>-</sup>		0.621 <sup>a</sup>	0.684 <sup>a</sup>	0.764 <sup>a</sup>	0.272 <sup>b</sup>
Ca <sup>2+</sup>			0.919 <sup>a</sup>	0.817 <sup>a</sup>	-0.408 <sup>a</sup>
Mg <sup>2+</sup>				0.811 <sup>a</sup>	-0.289 <sup>a</sup>
NH <sub>4</sub> <sup>+</sup>					-0.204
<i>Rancho Viejo, 1987–1993 (N = 267)</i>					
SO <sub>4</sub> <sup>2-</sup>	0.864 <sup>a</sup>	0.594 <sup>a</sup>	0.505 <sup>a</sup>	0.795 <sup>a</sup>	0.641 <sup>a</sup>
NO <sub>3</sub> <sup>-</sup>		0.660 <sup>a</sup>	0.518 <sup>a</sup>	0.828 <sup>a</sup>	0.602 <sup>a</sup>
Ca <sup>2+</sup>			0.744 <sup>a</sup>	0.587 <sup>a</sup>	0.188 <sup>a</sup>
Mg <sup>2+</sup>				0.490 <sup>a</sup>	0.062
NH <sub>4</sub> <sup>+</sup>					0.430 <sup>a</sup>

<sup>a</sup> Significant at *p* < 0.001.

<sup>b</sup> Significant at *p* < 0.05.

*N*: number of samples.

Table 3. Multiple regression analysis between H<sup>+</sup> and SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup> and NH<sub>4</sub><sup>+</sup> concentrations in Mexico City (1987–1990 and 1991–1993) and Rancho Viejo (1987–1993)

	Coefficient	<i>t</i> -value	Significant level	<i>F</i> -ratio ANOVA <sup>a</sup>	Significant level	<i>R</i>
<i>Mexico City, 1987–1990 (N = 104)</i>						
Constant	11.5166	3.0246	0.0032	64.2412	0.0000	0.748
Ca <sup>2+</sup>	-0.2689	-7.2256	0.0000			
NO <sub>3</sub> <sup>-</sup>	0.7368	11.2900	0.0000			
<i>Mexico City, 1991–1993 (N = 79)</i>						
Constant	-0.8666	-1.5239	0.1317	43.9656	0.0000	0.798
Ca <sup>2+</sup>	-1.8211	-6.5139	0.0000			
NH <sub>4</sub> <sup>+</sup>	-1.3530	-2.5254	0.0137			
SO <sub>4</sub> <sup>2-</sup>	3.6161	9.3308	0.0000			
<i>Rancho Viejo, 1987–1993 (N = 267)</i>						
Constant	-0.0869	-1.0114	0.3128	116.378	0.0000	0.684
SO <sub>4</sub> <sup>2-</sup>	0.9674	14.6707	0.0000			
Ca <sup>2+</sup>	-0.2545	-5.3508	0.0000			

<sup>a</sup> Analysis of variance for the full regression.

City is largely affected by dust particle emission sources. Magnesium and  $\text{SO}_4^{2-}$  only contribute 3%, and  $\text{NO}_3^-$  only 11%.

The above conclusions are strengthened by the results of chemical analyses performed on wet precipitation samples, also collected in Mexico City in 1991–1993. In this case, the  $\text{SO}_4^{2-}$  contribution increases to 5.5%. However,  $\text{Ca}^{2+}$  still has the strongest influence on  $\text{H}^+$  concentration (61%). Magnesium plus  $\text{NO}_3^-$  account for 2% and  $\text{NH}_4^+$  for 7%.

In summary, calcium is the most important ion in wet and bulk precipitation collected in Mexico City from 1991 to 1993 and from 1987 to 1990, respectively. This result strongly suggests that storm downdrafts blow off soil particles that are washed out by raindrops affecting their chemistry, and large soil particles are more rapidly scavenged during rainfall early stages (Tanaka *et al.*, 1980; Vong, 1990; Baez *et al.*, 1993; Camarero and Catalan, 1993). This explains why alkaline rainwater pHs are so common, despite the huge  $\text{SO}_2$  and  $\text{NO}_x$  emissions emitted in the Mexico City metropolitan area.

In this study carboxylic acids were not determined; however, one must consider the importance of their contribution in the acidity of precipitation.

Advances in sampling and analytical techniques have revealed that formic acid ( $\text{HCOOH}$ ) and acetic acid ( $\text{CH}_3\text{COOH}$ ) are major vapor (Dawson *et al.*, 1980) and aqueous (Keene *et al.*, 1983) phase constituents of the troposphere. Evidence based on direct measurements (Norton, 1985) and indirect estimates (Keene and Galloway, 1984) suggests that anthropogenic activities may represent important local sources for carboxylic acid distribution and origin. We believe that organic acids play an important role in precipitation acidity, since it is a well-known fact that Mexico City has one of the most photochemically polluted atmospheres in the world. Measurements of carboxylic acids are presently underway.

The ionic concentrations measured in rain water collected in Mexico City and Rancho Viejo can be compared with the results reported for other locations (Table 4). This table shows that sulfate concentrations measured at Rancho Viejo were the lowest. This result

Table 4. Comparison of wet and bulk precipitation ionic concentrations collected in two sites in Mexico with those reported in other locations

Site	$\text{SO}_4^{2-}$	$\text{NO}_3^-$	$\text{Ca}^{2+}$	$\text{Mg}^{2+}$	$\text{NH}_4^+$	$\text{H}^+$	References
	(μeq ℓ <sup>-1</sup> )						
<i>Wet precipitation</i>							
Minnesota, U.S.A. (seven sites average) (volume-weighted mean)	56.0	33.3	52.8	27.4	50.0	3.91	Pratt <i>et al.</i> (1983)
Warren, Michigan, U.S.A. (urban site, 42 weekly (average) data)	93.0	53.0	69.0	12.0	35.0	62.0	Dash and Cadle (1984)
South Western Europe (volume-weighted mean)	56.1	14.8	—	—	27.9	3.8	Pio <i>et al.</i> (1991)
Athens, Greece (year mean) (volume-weighted mean)	100	23.6	137	31.0	21.9	4.0	Dikaiakos <i>et al.</i> (1990)
Lower Ohio river, Kentucky, U.S.A. (all sites) (volume-weighted mean)	67.2	23.4	9.7	2.9	14.8	52.3	Saylor <i>et al.</i> (1992)
Mexico City, Mexico (all data) (volume-weighted mean)	75.3	34.8	33.1	3.7	76.1	22.3	This paper
Rancho Viejo, Mexico (all data) (volume-weighted mean)	39.7	15.6	9.3	5.5	26.9	20.5	This paper
<i>Bulk precipitation</i>							
Red Mountain Pass, Colorado (high altitude U.S.A. (geometric mean) sites, all data)	20.6	5.2	39.9	4.7	—	—	Reddy and Claassen (1985)
Greater Manchester, U.K. (volume-weighted mean)	85.4	29.3	49.5	26.5	31.4	43.0	Lee (1993)
Mexico City, Mexico (volume-weighted mean)	131	41.5	70.5	9.6	93.5	23.5	This paper



Table 5. Annual loading for rainfall collected at Mexico City and Rancho Viejo from 1987 to 1993

Year	SO <sub>4</sub> <sup>2-</sup>	NO <sub>3</sub> <sup>-</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	NH <sub>4</sub> <sup>+</sup>	H <sup>+</sup>	Rainfall (mm)
	g m <sup>-2</sup> yr <sup>-1</sup>						
<i>University of Mexico</i>							
1987	2.061	0.902	0.500	0.037	0.459	0.010	334.89
1988	2.080	0.680	0.449	0.036	0.582	0.004	306.93
1989	1.854	0.745	0.490	0.044	0.517	0.004	265.71
1990	1.545	0.755	0.248	0.024	0.455	0.009	288.58
1991	1.228	0.775	0.161	0.012	0.373	0.012	249.44
1992	1.626	0.802	0.262	0.018	0.521	0.0085	482.71
1993	0.895	0.663	0.263	0.017	0.415	0.0024	305.02
<i>Rancho Viejo</i>							
1987	0.266	0.127	0.038	0.007	0.064	0.0021	145.08
1988	0.417	0.193	0.068	0.012	0.096	0.0034	187.12
1989	0.639	0.303	0.058	0.013	0.150	0.0069	336.15
1990	0.584	0.290	0.035	0.006	0.153	0.0060	270.21
1991	0.526	0.254	0.037	0.005	0.117	0.0070	248.08
1992	0.352	0.179	0.029	0.003	0.085	0.0045	249.51
1993	0.381	0.262	0.043	0.005	0.139	0.0040	222.98

is expected since the sampling site is located in a rural wooded area with no near pollution sources and scarce road traffic; this fact is also valid for the NO<sub>3</sub><sup>-</sup> concentration measured there.

With respect to Mexico City, the SO<sub>4</sub><sup>2-</sup> levels in rain water are 1.46, 1.34 and 1.1 times higher than in Minnesota, South Western Europe and the Lower Ohio River, respectively, but only 0.75 and 0.80 times that in Athens, Greece and Warren, Michigan, respectively. It is interesting to note that the highest SO<sub>4</sub><sup>2-</sup> values correspond to urban areas affected by polluted air masses, like Warren (close to Detroit), Athens and Mexico City. Gatz (1980) has suggested that primary emissions may be the dominant source of SO<sub>4</sub><sup>2-</sup> in precipitation near or in large cities.

On the other hand, calcium is the ion that probably has the most important influence in rain acidity neutralization (Minnesota, 52.8 µeq ℓ<sup>-1</sup>; Warren 69 µeq ℓ<sup>-1</sup>), specially Athens (137 µeq ℓ<sup>-1</sup>) and Mexico City with values 0.62, 0.48, 0.24 times lower than in Minnesota, Warren and Athens, respectively.

The Ca<sup>2+</sup> concentration reported in these locations comes from different sources. In Minnesota, Pratt *et al.* (1983) reported that the sampling location may be the recipient of alkaline dust from the west. Calcium in Athens comes from soil dust (Dikaiakos *et al.*, 1990). Limestone is a predominant constituent of Greece's soils, and in Mexico City the Ca<sup>2+</sup> contribution to rain is from industrial particle alkaline emissions and soil erosion. It is very difficult to compare bulk precipitation results from other sites with those obtained in Mexico City, due mainly to the differences in sampling protocols between sampling locations. Reddy and Claassen (1985) collected bulk precipitation for seasonal periods, while Lee (1993) collected bulk samples on a weekly basis. The collectors were located in Greater Manchester, which covers an area of 1300 km<sup>2</sup> and with a wide range of activities such

as agriculture, industry, heavy traffic and a large power plant (2000 MW). In Mexico City, bulk precipitation was collected on a daily basis at the University of Mexico Campus 15 km south of Mexico City downtown and under the influence of the heavily polluted atmosphere of the city. Sulfate and Ca<sup>2+</sup> concentrations, 131.4 and 70.5 µeq ℓ<sup>-1</sup>, respectively, were higher than the values reported for Southwestern Europe and Greater Manchester. The high SO<sub>4</sub><sup>2-</sup> values found in Mexico City reflect the contribution of the SO<sub>2</sub> precursor that is emitted in huge quantities in Mexico City metropolitan area. The high calcium concentration seems to be due to the soil-derived compounds associated with wind-blown soil dust and from industrial alkaline particle emissions.

A summary of the rainy season annual deposition for rainfall collected in Mexico City and Rancho Viejo is presented in Table 5. As expected, higher deposition values correspond to bulk precipitation (Mexico City, from 1987 to 1990), since these values comprise dry and wet precipitation. The higher rainy season annual deposition in Mexico City possibly corresponds to high air pollution levels. In spite of the ionic concentration dilution by heavy rains, higher rainy season annual loading values correspond to higher rainfall amounts.

## CONCLUSIONS

A study of precipitation chemistry has been carried out for rain samples collected at two sampling sites in Central Mexico, over a 7-year period. The major conclusions are: The precipitation chemistry distribution was approximately log-normal but skewed toward higher values. Sulfate was the most abundant ion in all rain samples, followed by NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>. The monthly volume-weighted concentrations showed

that the highest ionic values were measured in May, when the rainy season usually begins in Central Mexico. The lowest concentrations were found from July to September, when most thunderstorms occur. The  $\log_{10}$  ion-pair correlations for the entire data set showed that the highest correlation corresponded to the ion pairs  $\text{Ca}^{2+}$ - $\text{Mg}^{2+}$ , which can be attributed to suspended soil dusts. The well-correlated pairs  $\text{SO}_4^{2-}$ - $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ - $\text{NH}_4^+$ , and  $\text{NO}_3^-$ - $\text{NH}_4^+$  seem to occur in precipitation because of common sources.

A multiple regression correlation analysis was performed after a  $\log_{10}$  transformation was applied to all data. The hierarchical model was used to select variables inserted into the final model. At Rancho Viejo,  $\text{SO}_4^{2-}$  was the largest contributor to the  $\text{H}^+$  concentration prediction. This result is consistent with the geographical characteristics and near-surface wind climatology of the sampling site. In contrast, in bulk precipitation collected in Mexico City,  $\text{Ca}^{2+}$  is the most important parameter, with a contribution of 38%, reflecting the influence of calcium compound emissions from industrial sources and wind-blown soil dust, in spite of high atmospheric  $\text{SO}_2$  levels. Even in wet precipitation collected in the city,  $\text{Ca}^{2+}$  still had the strongest influence on  $\text{H}^+$  ion concentration prediction. This fact suggests the strong influence of alkaline emissions and soil dust on  $\text{H}^+$  concentration.

Annual trends were not definitely discerned due to the high variability of the rainy season annual inputs of different ionic species.

The authors have considered the need to continue the study of precipitation chemistry in both Mexico City and a rural region downwind from the city, to determine whether the magnitudes of ionic inputs are increasing or decreasing and thus to assess present emission control effectiveness. In this research,  $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{K}^+$  and conductivity determinations will also be included to better understand the rain chemistry of the region.

*Acknowledgements*—The authors thank María del Carmen Torres B. and Rocío García for their help with the analytical work. We are also grateful to Dr Michel Douglas from NOAA for the review of the final draft of the manuscript and to the reviewers for their valuable suggestions.

## REFERENCES

- APHA (1985) *Standard Methods for the Examination of Water and Wastewater*. 16th Edn. APHA, AWWA, WPCF, Washington, District of Columbia.
- Asman H. A., Slanina J. and Board H. J. (1981) Meteorological interpretation of the chemical composition of rain-water at one measuring site. *Wat. Air Soil Pollut.* **16**, 161–175.
- Baez A. P. and Belmont R. (1987) Comparative study of the chemical composition of rain of three different zones in Mexico. *Contam. Ambient.* **3**, 25–36.
- Baez A. P., Belmont R. and Padilla G. H. (1993) Variation of chemical composition of wet precipitation, using a sequential sampling: urban–rural areas comparison. *Atmosfera* **6**, 163–74.
- Caiazza R., Hage D. K. and Gallup D. (1978) Wet and dry deposition of nutrient in Central Alberta. *Wat. Air Soil Pollut.* **9**, 309–314.
- Calvert J. G., Lazrus A., Kok G. L., Heikes B. G., Walega J. G., Lund J. and Cantrell C. A. (1985) Chemical mechanisms of acid generation in the troposphere. *Nature* **317**, 27–35.
- Camarero L. and Catalan J. (1993) Chemistry of bulk precipitation in the Central and Eastern Pyrenees, Northeast Spain. *Atmospheric Environment* **27A**, 83–94.
- Cohen J. and Cohen P. (1975) *Applied Multiple Regression, Correlation Analysis for the Behavioral Sciences*. Lawrence Erlbaum Associates, Hillsdale, New Jersey.
- Colin J. L., Jaffrezou J. L. and Gros J. M. (1990) Solubility of major species in precipitation: factors of variation. *Atmospheric Environment* **24A**, 537–544.
- Dash J. M. and Cadle S. H. (1984) The effect of local emissions on wet and dry deposition in southeastern Michigan. *Atmospheric Environment* **18**, 1009–1015.
- Daum P. H., Schwartz E. S. and Newman L. (1984) Acidic and related constituents in liquid water stratiform clouds. *J. geophys. Res.* **89**, 1147–1158.
- Dawson G. A., Farmer J. C. and Moyers J. L. (1980) Formic and acetic acids in the atmosphere of south-west U.S.A. *Geophys. Res. G. H.* **7**, 725–728.
- Dikaiakos J. G., Tsitouris C. G., Siskos P. A., Melissos D. A. and Nastos P. (1990) Rain water composition in Athens, Greece. *Atmospheric Environment* **24B**, 171–176.
- Galloway J. N. and Likens G. E. (1978) The collection of precipitation for chemical analysis. *Tellus* **30**, 71–82.
- Galloway J. N., Likens G. E. and Hawley M. E. (1984) Acid precipitation: natural versus anthropogenic components. *Science* **226**, 829–831.
- Gatz D. F. (1980) Association and mesoscale relationships among rain water constituents. *J. geophys. Res.* **85**, 5588–5598.
- Jáuregui E. (1986) The urban climate of Mexico. In *Proc. WMO Tech. Conf. on Urban Climate in Tropical Areas*. (edited by Oke T. R.), WMO 652, pp. 63–83.
- Keene W. C. and Galloway J. N. (1984) Organic acidity in precipitation of North America. *Atmospheric Environment* **18**, 2491–2497.
- Keene W. C., Galloway J. N. and Holden J. D. Jr (1983) Measurement of weak organic acidity in precipitation from remote areas of the world. *J. geophys. Res.* **88**, 5122–5130.
- Kins L. (1982) Temporal variation of chemical composition of rain water during precipitation events. In *Deposition of Atmospheric Pollutants* (edited by Georgii H. W. and Pankrath J.), pp. 87–96. Reidel, Dordrecht.
- Lee D. S. (1993) Spatial variability of urban precipitation chemistry and deposition: statistical associations between constituents and potential removal processes of precursor species. *Atmospheric Environment* **27B**, 321–337.
- Mészáros E. (1978) *Background Air Pollution Monitoring*. Institute for Atmospheric Physics of the Meteorological Service of Hungary. Prepared for the World Meteorological Organization and The United Nations Environment Programme Training Course.
- Norton R. B. (1985) Measurements of formate and acetate in precipitation at Niwot Ridge and Boulder, Colorado. *J. geophys. Res.* **12**, 769–772.
- Padilla H. G., Leyva A. C. and Mosiño P. A. (1993) An analysis of daily humidity patterns at a mountainous and urban site in a tropical high-altitude region. *J. appl. Met.* **32**, 1638–1646.
- Parungo F., Nagamoto C., Hoyt S. and Bravo H. A. (1990) The investigation of air quality and acid rain over the Gulf of Mexico. *Atmospheric Environment* **24A**, 109–123.
- Peden M. E. and Skowron L. M. (1978) Ionic stability of precipitation samples. *Atmospheric Environment* **12**, 2343–2349.

- Penkett S. A., Jones B. M. R., Brice K. A. and Eggleton E. J. (1979) The importance of atmospheric ozone and hydrogen peroxide in oxidizing sulfur dioxide in cloud and rain water. *Atmospheric Environment* **13**, 123–137.
- Pio C. A., Salgueiro M. L. and Nunes T. V. (1991) Seasonal and air-mass trajectory effects on rainwater quality at the South-western European Border. *Atmospheric Environment* **25A**, 2259–2266.
- Prado-Fiedler R. (1990) On the relationship between precipitation amount and wet deposition of nitrate and ammonium. *Atmospheric Environment* **24A**, 3061–3065.
- Pratt G. C., Coscio M., Gardner D. W., Chevone B. I. and Krupa S. V. (1983) An analysis of the chemical properties of rain in Minnesota. *Atmospheric Environment* **17**, 347–355.
- Reddy M. M. and Claassen H. C. (1985) Estimates of average major ion concentrations in bulk precipitation at two high-altitude sites near the Continental Divide in South-western Colorado. *Atmospheric Environment* **19**, 1199–1203.
- Saylor R. D., Butt K. M. and Peters L. K. (1992) Chemical characterization of precipitation from a monitoring network in the lower Ohio River Valley. *Atmospheric Environment* **26A**, 1147–1156.
- Schwartz S. E. (1984) Gas–aqueous-reactions of sulfur and nitrogen oxides in liquid-water clouds. In *SO<sub>2</sub>, NO and NO<sub>2</sub> Oxidation Mechanisms: Atmospheric Considerations* (edited by J. G. Calvert), pp. 173–208. Butterworths, Boston.
- Seinfeld J. H. (1986) *Atmospheric Chemistry and Physics of Air Pollution*. Wiley, New York.
- Servicio Meteorológico Nacional (1992) Personal communication.
- Tanaka S., Darzi M. and Winchester J. W. (1980) Short term effect of rainfall on elemental composition and size distribution of aerosols in North Florida. *Atmospheric Environment* **14**, 1421–1426.
- Vong R. J. (1990) Mid-latitude northern hemisphere background sulfate concentration in rain water. *Atmospheric Environment* **24A**, 1007–1018.